



Costantini, D. (2016) Oxidative stress ecology and the d-ROMs test: facts, misfacts and an appraisal of a decade's work. *Behavioral Ecology and Sociobiology*, 70(5), pp. 809-820.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/120076/>

Deposited on: 14 July 2016

Enlighten – Research publications by members of the University of Glasgow  
<http://eprints.gla.ac.uk>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15

**Oxidative stress ecology and the d-ROMs test:  
facts, misfacts and an appraisal of a decade's work**

David Costantini

Department of Biology, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium  
Institute of Biodiversity, Animal Health and Comparative Medicine, School of Life  
Sciences, University of Glasgow, Glasgow G12 8QQ, UK

Email: [davidcostantini@libero.it](mailto:davidcostantini@libero.it)

**Running head:** Oxidative stress and the d-ROMs test

16 **Abstract** In recent years, behavioural ecologists have taken to studying oxidative stress in  
17 free-ranging organisms because it has been proposed as an important mediator of life-  
18 history trade-offs. A plethora of methodological approaches to quantify biomarkers  
19 associated with oxidative stress exist, each one with its own strengths and weaknesses. The  
20 d-ROMs test has emerged as one of the favoured assays in ecological studies because of its  
21 reliability, sensitivity to specific perturbations of the organism's oxidative balance, and  
22 medical and ecological relevance. Criticisms have been, however, raised about its  
23 specificity for oxidative damage. In this article, I have reviewed basic information about the  
24 d-ROMs test, its validation, the methodological mistakes made in the studies that attempted  
25 to criticise this assay and the application of this assay to ecological studies of oxidative  
26 stress. All the direct and indirect evidence shows that the d-ROMs test is a valuable assay  
27 for the quantification of plasma or serum primary (or early) oxidative damage molecules  
28 and, possibly, of other biological matrices and provides ecologically relevant information.

29

30 **Keywords** Biomarker • Life-history • Oxidative damage • Physiology

31

## 32 **Introduction**

33 Oxidative stress is a complex multifaceted biochemical condition of cells, which occurs  
34 when there is an increased rate of oxidative molecular damage and oxidation of non-protein  
35 and protein thiols that regulate the cell oxidative balance (Sies 1991; Jones 2006; Halliwell  
36 and Gutteridge 2007). Although biomedical and toxicological sciences have traditionally  
37 centralised research on oxidative stress, in recent years, ecologists have also taken to  
38 studying oxidative stress in free-ranging organisms and have integrated principles of  
39 oxidative stress into several core evolutionary concepts, such as life-history trade-offs (e.g.  
40 survival vs. reproduction), senescence and sexual selection. It is increasingly recognised  
41 that the need to manage the oxidative status in an optimal way may be an important  
42 mechanism driving the outcome of many life-history trade-offs (Costantini 2008, 2014;  
43 Metcalfe and Alonso-Alvarez 2010; Blount et al. 2015; Tobler et al. 2015).

44         There are many methodological approaches for the assessment of different  
45 biomarkers of oxidative status, including assays of oxidative damage, enzymatic or non-  
46 enzymatic antioxidants and repair molecules (Dotan et al. 2004; Halliwell and Gutteridge  
47 2007). However, there is no single biochemical metric that fulfils the need to properly  
48 quantify the organism oxidative balance (Dotan et al. 2004). It has therefore been  
49 repeatedly recommended to couple experimental manipulations with comprehensive  
50 metrics of oxidative status. In ecological research, there are also specific restrictions  
51 inherent to many research programmes, such as the availability of only small amounts of  
52 blood and the requirement of non-terminal sampling. Hence, ecologists have been mostly  
53 relying on those biomarkers of oxidative status that can be measured in blood (e.g., plasma,  
54 red blood cells). Much work has involved markers of oxidative damage, including end-

55 products of lipid peroxidation (malondialdehyde), damage to proteins (protein carbonyls) or  
56 products of oxidative damage that are generated early in the oxidative cascade (organic  
57 hydroperoxides). The d-ROMs test has emerged as one of the favoured assays for the  
58 quantification of some aspects of the plasma oxidative status in ecological studies. The d-  
59 ROMs has enabled to characterise many significant associations between plasma oxidative  
60 status and either physiological or life-history traits (Table 1). However, some mistakes have  
61 been made in the interpretation of what the d-ROMs actually measures. Moreover, a few  
62 criticisms have been raised about the specificity of the d-ROMs. Hence, in this article, I  
63 have reviewed basic information about organic hydroperoxides because these are the main  
64 molecules measured by the d-ROMs; the technical aspects of the d-ROMs; the validation of  
65 this assay; the methodological mistakes made in the studies that attempted to criticise this  
66 assay; and the application of this assay to ecological studies of oxidative stress.

67

### 68 **What are organic hydroperoxides?**

69 Organic hydroperoxides derive from the oxidation of biomolecular substrates, such as  
70 polyunsaturated fatty acids, cholesterol, proteins and nucleic acids, and are precursors of  
71 end-products of lipid peroxidation, such as malondialdehyde, hydroxynonenal and  
72 isoprostanes (Halliwell and Gutteridge 2007; Lajtha et al. 2009). Organic hydroperoxides  
73 are therefore biomolecules that were damaged by free radicals and, as such, lost their  
74 functionality. In plasma of vertebrates, baseline organic hydroperoxides concentrations are  
75 usually below 10  $\mu$ M (e.g., examples on birds and mammals in Miyazawa 1989; Gerardi et  
76 al. 2002; Montgomery et al. 2011, 2012).

77           An important source of organic hydroperoxides is peroxidation of lipids. Reactive  
78 species are capable of abstracting a hydrogen atom from polyunsaturated fatty acids (Fig.  
79 1), thus initiating a chain reaction known as lipid peroxidation. During this process,  
80 membrane lipids are oxidized yielding lipid organic hydroperoxides as primary products.  
81 Cells are equipped with enzymes belonging to the glutathione peroxidase family capable of  
82 reducing organic hydroperoxides to less toxic molecules. However, organic hydroperoxides  
83 can also accumulate to some extent and participate in reactions that fuel oxidative stress  
84 and increase toxicity. For example, the toxicity of organic hydroperoxides is promoted by  
85 the presence of metals which catalyse their cleavage (Fenton reaction), leading to the  
86 generation of two highly reactive and histolesive pro-oxidants, namely the alkoxyl (R-O•)  
87 and alkylperoxyl (R-OO•) radicals (Girotti 1998). Organic hydroperoxides are known to  
88 alter cell membrane fluidity and properties and to promote cell necrosis and death (Kagan  
89 et al. 2004). A wide range of organic hydroperoxides can also be formed from the reaction  
90 of proteins (Fig. 2) or nucleic acids (Fig. 3) with reactive species. For example, amino acids  
91 like proline, glutamate or lysine have been found to generate hydroperoxides (Simpson et  
92 al. 1992). Also, it has been found that reaction between hydroxyl radical and nucleic acids  
93 can generate several DNA hydroperoxides, such as the hydroxy-6-hydroperoxy-5,6-  
94 dihydrothymidine or the 6-hydroxy-5-hydroperoxide-5,6-dihydro-2'-deoxycytidine (Cadet  
95 and Di Mascio 2006; Miyamoto et al. 2007).

96

### 97 **Assays for measuring organic hydroperoxides**

98 There are several methodological approaches for the quantification of organic  
99 hydroperoxides in serum and plasma or in other biological matrices. These methods include



100 (i) colorimetric assays (e.g., FOX2 assay, d-ROMs assay, LPO assay kit) and flow  
101 cytometry (Sirak et al. 1991) for the quantification of total or lipid hydroperoxides and (ii)  
102 chromatographic techniques (e.g., gas chromatography-mass spectrometry, Kulmacz et al.  
103 1990; high performance liquid chromatography with chemiluminescence detection, Ferretti  
104 et al. 2005) for the specific quantification of certain groups of organic hydroperoxides. Of  
105 these methods, the d-ROMs assay has been increasingly used because of its high  
106 performance in terms of intra- and inter-coefficient of variation (below 10%), recovery rate  
107 (e.g., between 92 and 106% in Vassalle et al. 2006 and Pasquini et al. 2008) and linearity of  
108 dilution series of plasma samples (e.g.,  $R^2 \geq 0.95$  in Pasquini et al. 2008).

109


#### 110 **Description of the d-ROMs test**

111 The d-ROMs test measures the oxidant ability of a serum or plasma sample towards a  
112 particular substance (modified aromatic amine) used as an indicator (chromogen). Organic  
113 hydroperoxides are the compounds that mainly contribute to such oxidant ability, hence  
114 providing an indirect estimate of organic hydroperoxides (Alberti et al. 2000). The  
115 quantification of organic hydroperoxides is indirect because the acidic pH of the buffer  
116 used for the reaction favours the release of metal ions like iron ( $Fe^{2+}$  and  $Fe^{3+}$ ) and copper  
117 ( $Cu^+$  and  $Cu^{2+}$ ) from circulating proteins. These metals catalyse the cleavage of organic  
118 hydroperoxides, leading to the generation of two free radicals, i.e.,  $RO\bullet$  and  $ROO\bullet$ , that  
119 oxidize the alkylated amine DEPPD (*N,N*-diethyl-*para*-phenyldiamine) contained in the  
120 chromogen solution to its red radical cation, the intensity of the resulting colour being  
121 related to the amount of organic hydroperoxides present in the sample. The overall amount  
122 of metal ions that can trigger the Fenton reaction occurs in excess in plasma or serum (i.e.,

123 > 10  $\mu\text{M}$ ; e.g., Lumeij and de Bruijne 1985; Suominen et al. 1988; Spolders et al. 2010;  
124 Kautz et al. 2014) as compared to the amount of organic hydroperoxides, hence it is  
125 unlikely that the availability of metal ions might limit the performance of the assay.  
126 Moreover, according to the following reactions, both iron and copper ions are reduced back  
127 continuously after they were oxidised, avoiding any reduction in their availability:



131 and



134 The results of the d-ROMs test are expressed in arbitrary units called “Carratelli units”  
135 (CARR U), where 1 CARR U is equivalent to 0.08 mg of  $\text{H}_2\text{O}_2/100 \text{ mL}$  or to 0.024mM  
136  $\text{H}_2\text{O}_2$  equivalents. Note that, for practical reasons, the value is expressed as a chemical  
137 equivalence and not as a concentration. This is also because as stated in the early study  
138 where the d-ROMs assay was described (Alberti et al. 2000), as well as in next studies  
139 (Liang et al. 2012), while organic hydroperoxides are the main molecules detected by the d-  
140 ROMs assay, other primary oxidative damage molecules may also be detected (e.g., organic  
141 chloramines that derive from oxidation of protein amine groups; endoperoxides in Liang et  
142 al. 2012). According to the Marcus theory (1956), chemical species with a reduction

143 potential higher than 0.3 V toward a standard hydrogen electrode should theoretically be  
144 able to oxidise the DEPPD, but their contribution might be relevant as long as they occur at  
145 a concentration comparable to that of organic hydroperoxides.

146 Electron paramagnetic resonance (EPR) spectroscopy is the gold standard technique  
147 to detect the formation of free radicals in a biological matrix and to identify their nature. In  
148 an early EPR investigation of the d-ROMs test (Alberti et al. 2000), it was possible to  
149 unambiguously identify the radical cation DEPPD<sup>•+</sup> as the species responsible for the  
150 colour formation and to show that the free radicals originating from the cleavage of organic  
151 hydroperoxides did lead to the formation of DEPPD<sup>•+</sup>. This early assessment of the d-  
152 ROMs test was replicated by another EPR investigation, which supported the conclusion  
153 that oxidants like organic hydroperoxides are the main molecules that are quantified by the  
154 d-ROMs test (Liang et al. 2012). Liang et al. (2012) also suggested that other oxidative  
155 damage molecules like endoperoxides contribute to the d-ROMs values.

156 It has been argued that ceruloplasmin (glycoprotein produced in the liver that carries  
157 copper in blood plasma) may have some interference in the d-ROMs readings due to its  
158 ferroxidase activity and to the ability of its Cu<sup>2+</sup> ions to oxidize the amine DEPPD (e.g.,  
159 Erel 2005; Harma et al. 2006). Another possibility is that ceruloplasmin can indirectly  
160 contribute to the detection of organic hydroperoxides because this protein exerts a  
161 ferroxidase activity by converting ferrous to ferric ions and the resulting Cu<sup>+</sup> ions are in  
162 turn amenable to react with hydroperoxides to give alkoxyl radicals in a Fenton-like  
163 reaction (Colombini et al. 2016). However, experiments with a ceruloplasmin inhibitor  
164 (NaN<sub>3</sub>) added in excess as compared to the ceruloplasmin concentration suggested that  
165 ceruloplasmin only plays a minor role (around 7%) in determining the d-ROMs test

166 readings in human sera (Alberti et al. 2000) and even lower in other species (Table 2).  
167 Although the amount of sodium azide used by Alberti et al. (2000) was chosen without  
168 consideration of the affinity of sodium azide for ceruloplasmin, inhibition of ceruloplasmin  
169 with another molecule (the copper chelator bathocuproine disulfonate) confirmed the  
170 minimal interference of ceruloplasmin (Liang et al. 2012). If the d-ROMs assay detects  
171 only ceruloplasmin as suggested by criticisms, it should be expected to find a very high  $R^2$ ,  
172 probably higher than 0.90, between d-ROMs values and ceruloplasmin. New recent studies  
173 found a weak correlation between the results of the d-ROMs assay and the amount of  
174 ceruloplasmin actually present in the samples ( $R^2 = 0.0009$ , Costantini et al. 2014a;  $R^2 =$   
175  $0.18$  or  $R^2 = 0.07$  when one outlier outside the 99% confidence interval is excluded,  
176 Colombini et al. 2016). Moreover, Colombini et al. (2016) showed that the addition of  
177 ceruloplasmin (2.27 or 6.8  $\mu\text{M}$ ) to human serum samples caused on average an increase of  
178 around 6% of the d-ROMs values, which is in agreement with previous work.

179 In Buonocore et al. (2000) several determinations, adding purified glutathione  
180 peroxidase with reduced glutathione in excess to the plasma sample, were performed to  
181 prove that the reaction of the d-ROMs test was due to peroxy and alkoxy radical products.  
182 This procedure was used because glutathione is used by the enzyme glutathione peroxidase  
183 to reduce organic hydroperoxides to their corresponding alcohols. In all determinations  
184 made, a decrease of more than 90% in the signal was observed, supporting early work by  
185 Alberti et al. (2000). Although glutathione is per se a scavenger of free radicals (i.e., it  
186 provides its hydrogen atom to radical anions), the amine used by the d-ROMs assay is a  
187 radical cation, hence it cannot accept the hydrogen atom from glutathione. Hence, a direct  
188 reaction between glutathione and the amine can be excluded. This is further demonstrated

189 by a recent study, where experimental decrease of glutathione by administration of  
190 sulfoximine was associated with increased d-ROMs values (Costantini et al. 2016; Fig. 4).  
191 Although glutathione and d-ROMs were measured in red blood cells and plasma,  
192 respectively, note that it is well established that sulfoximine reduces glutathione synthesis  
193 in the whole body, including kidney (Griffith and Meister 1979), jejunum, lung, heart, liver  
194 and brain (Favilli et al. 1997) and plasma (Ovrebø et al. 1997; Ovrebø and Svardal 2000).  
195 Finally, note that Kilk et al. (2014) found that glutathione did not affect the d-ROMs values  
196 at normal serum values.

197

#### 198 **Indirect evidence: sensitivity of the d-ROMs assay to pro-oxidant agents**

199 A way to test the sensitivity of an assay to perturbations of the organism's oxidative  
200 balance is to manipulate the free radical production of an organism using methods that  
201 specifically increase basal production of free radicals, such as exposure to hyperoxia or  
202 injection of generators of reactive species or of inhibitors of antioxidants. Several  
203 experimental reports have clearly shown that the d-ROMs test is very sensitive to specific  
204 manipulations of oxidative stress level. For example, during 5 and 15 minutes of  
205 reperfusion (i.e., restoration of the flow of blood to a previously ischemic tissue or organ)  
206 of Syrian hamsters, plasma values of d-ROMs increased by 72% and 89%, respectively, as  
207 compared to baseline values and declined to baseline after 30 minutes of reperfusion  
208 (Bertuglia and Giusti 2003). However, pretreatment of Syrian hamsters with the antioxidant  
209 enzyme superoxide dismutase maintained d-ROMs values at normal levels, indirectly  
210 showing that most molecules detected by the d-ROMs test come from oxidation of  
211 biomolecules induced by free radicals. Similarly, Benedetti et al. (2004) found that the

212 repeated exposures of patients to hyperbaric oxygen (i.e., 100% oxygen) led to a significant  
213 increase in plasma d-ROMs values, as well as in another biomarker of oxidative damage  
214 (thiobarbituric acid reactive substances). In another similar experiment, Nagatomo et al.  
215 (2012) found that there were no differences in d-ROMs values in rats exposed to 14.4%,  
216 20.9%, and 35.5% oxygen. However, d-ROMs values increased in the rats exposed to  
217 39.8% and 62.5% oxygen. d-ROMs values were the highest in the rats exposed to 82.2%  
218 oxygen. Morphological changes in the red blood cells induced by oxidative attack from  
219 reactive oxygen species were also observed in the rats exposed to 39.8%, 62.5%, and  
220 82.2% oxygen (Nagatomo et al. 2012). In another experiment, domestic canaries were  
221 injected with DL-buthionine-(S,R)-sulfoximine, a compound that reduces cellular levels of  
222 glutathione by inhibiting its synthesis (Griffith and Meister 1979; Bailey 1998). As  
223 compared to controls, canaries treated with sulfoximine had a significant decrease of  
224 reduced glutathione and a significant increase of d-ROMs values, respectively; in contrast,  
225 the activity of ceruloplasmin was not affected by the treatment (Costantini et al. 2016; Fig.  
226 4).

227

#### 228 **Further considerations on the d-ROMs assay**

229 Among species variation in ceruloplasmin and d-ROMs values also provides indirect  
230 support for the specificity of the d-ROMs assay. For example, while humans and pigs have  
231 similar activity of ceruloplasmin (Schosinsky et al. 1974; Feng et al. 2007), d-ROMs values  
232 in pigs are 2 to 6 times higher than in humans (humans, 325 UC  $\approx$  7.8 mM H<sub>2</sub>O<sub>2</sub>  
233 equivalents in Schöttker et al. 2015b; pigs, 558 to 1750 UC  $\approx$  13.4 to 42.0 mM H<sub>2</sub>O<sub>2</sub>  
234 equivalents in Brambilla et al. 2002). Similarly, while ceruloplasmin is already active and

235 detectable in young individuals at levels similar to those of adults (Lin et al. 2004; Fleming  
236 et al. 2009; Wang et al. 2014), d-ROMs values may not be detectable in plasma of young  
237 Eurasian kestrel individuals or are lower than those of adults (young kestrels in Costantini  
238 et al. 2006; adult kestrels in Casagrande et al. 2011; Vassalle et al. 2006 for an example on  
239 humans; Sgorbini et al. 2015 for an example on horses).

240         Although, in general, the patterns of variation of d-ROMs values and ceruloplasmin  
241 differ between each other implying low or no correlation between them (see e.g., Assenza  
242 et al. 2009; Talukder et al. 2014, 2015), it has to be considered that it is not correct to infer  
243 about a potential interference of a given molecule in an assay from a simple correlation.  
244 Ceruloplasmin contributes to the regulation of oxidative balance in the organism (Goldstein  
245 et al. 1979; Calabrese and Carbonaro 1986; Samokyszyn et al. 1989; Ehrenwald et al.  
246 1994), hence it can happen to find or not to find correlations between ceruloplasmin and  
247 markers of oxidative damage, including organic hydroperoxides (Bednarek et al. 2004;  
248 Maykova et al. 2013; Saravanan and Ponmurugan 2013; Kusuma Kumari and  
249 Sankaranarayana 2014).

250         Results obtained with the d-ROMs test are also consistent with studies that  
251 quantified organic hydroperoxides using other methods. For example, increased production  
252 of organic hydroperoxides (as measured by the d-ROMs) during an immune/inflammatory  
253 response is in agreement with other studies that quantified the production of organic  
254 hydroperoxides using different methods (e.g., effect of 12-O-tetradecanoylphorbol-13-  
255 acetate on organic hydroperoxides in Sirak et al. 1991; effect of lipopolysaccharide on  
256 organic hydroperoxides in Riedel et al. 2003; effect of multiple sclerosis on organic  
257 hydroperoxides in Ferretti et al. 2005). This strong and significant link between organic

258 hydroperoxides and immune response is further corroborated by the significant correlations  
259 that have been found between d-ROMs values and leucocyte counts or antibody titres (e.g.,  
260 van de Crommenacker 2011; Casagrande et al. 2012; Schneeberger et al. 2013). Note also  
261 that injection of pigeons with *Escherichia coli* lipopolysaccharide caused inflammation, but  
262 did not induce any relevant changes in ceruloplasmin (Dudek et al. 2013). Increased  
263 production of organic hydroperoxides was also found in relation to increased levels of  
264 stress hormones using either the d-ROMs (Costantini et al. 2008; Haussmann et al. 2012) or  
265 other assays (Sato et al. 2010; Balkaya et al. 2011).

266         Some authors used the d-ROMs assay to estimate production of free radicals (Al-  
267 Johany et al. 2009; Noguera et al. 2011). Organic hydroperoxides are likely to better reflect  
268 the basal free radical production than endproducts of lipid peroxidation (e.g.,  
269 malondialdehyde). This is because organic hydroperoxides are generated earlier in the  
270 oxidative cascade than endproducts. However, organic hydroperoxides are primary  
271 oxidative damage molecules and none studies have tested the correlation between organic  
272 hydroperoxides and amount of free radicals produced. The correlation between free radical  
273 generation and organic hydroperoxides is unlikely to be very strong because there are  
274 various mechanisms and molecules that either prevent oxidation of biomolecules or remove  
275 organic hydroperoxides from the organism. It is therefore premature to infer about the  
276 production of free radicals from the results of the d-ROMs assay.

277         It has also been suggested that the d-ROMs assay is a measure of potential damage  
278 (Stier et al. 2015). This sentence implies that the d-ROMs test detects molecules that may  
279 potentially cause damage. This may certainly happen because molecules like organic  
280 hydroperoxides can be cleaved into free radicals, so fuelling the oxidative cascade.

281 However, it is unknown the extent to which this can happen (e.g., peroxidases reduce  
282 organic hydroperoxides to their corresponding alcohols before they can be cleaved). It  
283 therefore appears more correct to refer to oxidative damage because this is what the main  
284 molecules detected by the d-ROMs assay are. Note that, although peroxidases interact with  
285 organic hydroperoxides in the organism, it is unlikely that peroxidases cause significant  
286 interference with the d-ROMs test because of the low activity of peroxidases in plasma and  
287 the weak correlation between peroxidase activity and d-ROMs values (Costantini et al.  
288 2011, 2012a).

289

#### 290 **Beyond the blood: application of the d-ROMs to other biological matrices**

291 Although the application of the d-ROMs test has been originally assessed for blood, recent  
292 studies showed that the d-ROMs test might also be applied to other biological matrices. For  
293 example, Castellini et al. (2003) found that d-ROMs values of seminal plasma of rabbits are  
294 correlated to those measured in blood plasma. Ito et al. (2009) found that plasma d-ROMs  
295 values were positively and significantly correlated ( $r = 0.50$ ) with urine d-ROMs values. In  
296 another study, the d-ROMs test was applied to immune cells. Specifically, bovine  
297 peripheral blood mononuclear cells (component of the immune system) had been isolated  
298 from the whole blood and were then incubated for 2 and 7 days at different concentrations  
299 of mycotoxins to stimulate their activity (Bernabucci et al. 2011). Compared with the  
300 control, an increase of intracellular d-ROMs values (and also of malondialdehyde, which is  
301 an endproduct of lipid peroxidation) was observed, indicating the high sensitivity of the d-  
302 ROMs assay to perturbations of the oxidative balance caused by an immune response.

303 Similarly, in vitro induction of oxidative stress in prostate cancer cells resulted in  
304 the increase of d-ROMs values in the extracellular compartment, and of reactive oxygen  
305 species and DNA damage in the intracellular compartment (Tomasetti et al. 2010).

306 Finally, Criscuolo et al. (2010) found that the d-ROMs test may also be applied to  
307 haemolymph of invertebrates. Criscuolo et al. (2010) found that the d-ROMs values were  
308 higher in the haemolymph of short-lived male tarantulas (which also had higher superoxide  
309 production and lower antioxidant defences) than in the haemolymph of their long-lived  
310 females (which also had lower superoxide production and higher antioxidant defences).

311

#### 312 **Does the d-ROMs assay provide ecologically relevant information?**

313 Medical and veterinary research showed that the d-ROMs test has a significant diagnostic  
314 value of many pathological statuses, in agreement with other biomarkers of oxidative  
315 damage, such as the gold standard “isoprostanes” (e.g., correlation between d-ROMs values  
316 and isoprostanes = 0.68 in Lubrano et al. 2002). For example, d-ROMs values were found  
317 to be significantly associated with mortality in humans independently from established risk  
318 factors, including inflammation (Schöttker et al. 2015a).

319 It is now increasingly recognised that the d-ROMs is also a valuable test for the  
320 investigation of oxidative stress in ecological studies. Work has, for example, shown that d-  
321 ROMs values are associated with behavioural or fitness-related traits or reflect potential  
322 physiological costs induced by short-term activation of immune response or changes in the  
323 hormonal profile (Table 1). For example, experimental increase of plasma d-ROMs values  
324 was found to delay egg laying and reduce clutch size in a songbird (Costantini et al. 2016),  
325 which are two important fitness-related traits under female control. High plasma d-ROMs

326 values were found to be associated with reduced survival perspectives in two seabird  
327 species (Costantini et al. 2015; Herborn et al. 2016). Geiger et al. (2012) and Hau et al.  
328 (2015) also found that high plasma d-ROMs values were associated with shorter telomeres,  
329 which is an emerging marker of disease risk and biological ageing. Finally, a number of  
330 studies found that the d-ROMs test may inform about environmental quality (e.g.,  
331 contamination level in Bonisoli-Alquati et al. 2010; food availability in van de  
332 Crommenacker et al. 2011a), adaptation to urbanisation (Lucas and French 2012;  
333 Costantini et al. 2014b) or variation in behavioural phenotype (Herborn et al. 2011;  
334 Costantini et al. 2012b).

335

#### 336 **Criticisms on the d-ROMs assay are based on methodological inadequacy**

337 A few authors raised critics to the d-ROMs assay, claiming that it is not a valid assay for  
338 the quantification of plasma oxidative status, organic hydroperoxides in particular.  
339 However, these critics have been based on serious methodological inadequacy and,  
340 importantly, on lack of respect of the protocols that have previously been validated.

341 Early criticisms about the lack of specificity were based on several mistakes done in  
342 the application of the d-ROMs assay (Erel 2005; Harma et al. 2006). For example, Erel  
343 (2005) used chelants for the preparation of samples, which interfere with the d-ROMs  
344 reaction (Banfi et al. 2006). Chelants sequester iron ions, which are needed for the reaction  
345 of the d-ROMs test and this was already explained in the validation study (Alberti et al.  
346 2000). Hence, anticoagulants like EDTA or citrate should not be used.

347 Recently, similar critics to those of Harma et al. (2006) were surprisingly raised by  
348 Kilk et al. (2014). Again, these critics have been based on methodological inadequacy in

349 the application of the d-ROMs assay. First, Kilk et al. (2014) tested the capacity of the d-  
350 ROMs assay to assess oxidative damage using solutions of hydrogen peroxide. They found  
351 that the d-ROMs was able to weakly detect hydrogen peroxide only when it occurred at  
352 very high concentrations ( $\geq 50 \mu\text{M}$ ) that are sometimes found in the plasma of individuals  
353 under severe oxidative stress. Hence, the authors concluded that the amount of peroxides in  
354 the plasma is a bit above the detection limit of the d-ROMs assay only under conditions of  
355 strong oxidative stress. The small capacity of the d-ROMs to measure hydrogen peroxide  
356 is, however, not surprising nor is it novel (see Liang et al. 2012) because the d-ROMs  
357 mainly measures organic hydroperoxides (e.g., Alberti et al. 2000; Liang et al. 2012;  
358 Colombini et al. 2016). The d-ROMs can detect organic hydroperoxides at concentrations  
359 well below  $10 \mu\text{M}$  (Fig. 5), which have been found in the plasma of several vertebrates  
360 (e.g., Miyazawa 1989; Gerardi et al. 2002; Montgomery et al. 2011, 2012). For example,  
361 Gerardi et al. (2002) found a significant positive correlation between d-ROMs values and  
362 lipid hydroperoxides measured with the FOX2 assay.

363         Second, Kilk et al. (2014) found that d-ROMs readings of solutions containing  
364 ceruloplasmin and of sera decreased by decreasing the incubation temperature from  $37^\circ\text{C}$   
365 (protocol of d-ROMs) to  $23^\circ\text{C}$  and to  $4^\circ\text{C}$ , while those of solutions containing hydrogen  
366 peroxide were less dependent on temperature. These results are again unreliable because (i)  
367 the d-ROMs assay is poorly sensitive to hydrogen peroxide, (ii) the Fenton reaction is  
368 highly dependent on temperature (e.g., Neyens and Baeyens 2003; Lee and Yoon 2004;  
369 Hussain et al. 2014) as also shown by the decrease of the readings of pure solutions of  
370 cumene hydroperoxide with the decrease of incubation temperature (Table 3), and (iii) the  
371 decrease of absorbance of pure solutions of cumene hydroperoxide with temperature is

372 similar to that of plasma samples (Table 3; note that the pure solution of organic  
373 hydroperoxides has to have an absorbance at 37°C similar to that of plasma samples in  
374 order to avoid “the regression to the mean effect”, i.e., when the change in absorbance is  
375 dependent on the initial absorbance), while that of ceruloplasmin shown in Kilk et al.  
376 (2014) is not.

377 Third, the authors speculated that thiols like albumin and glutathione might interfere  
378 with the assay. However, the amount of either albumin or glutathione that Kilk et al. (2014)  
379 added to the serum samples was pharmacological, while, as stressed by the same authors,  
380 glutathione did not affect the d-ROMs values at normal serum values.

381 Fourth, given this apparent temperature effect, the authors stated that they carried  
382 out the incubation at 23°C. Although the early assessment of the d-ROMs performance was  
383 carried out at room temperature (Alberti et al. 2000), the decrease of Fenton reaction with  
384 incubation temperature shown in Table 3 suggests that several of the conclusions made by  
385 the authors should be taken cautiously.

386 Fifth, the supposed correlation between d-ROMs values and activity of  
387 ceruloplasmin shown in figure 7 is not reliable because (i) it is an artifact of plotting two  
388 species having different levels of both d-ROMs values and ceruloplasmin and (ii) more  
389 importantly because the values of absorbance of sera reported in figure 7 are not compatible  
390 with the normal absorbance values of the assay (see e.g. values of absorbance in Table 2 for  
391 a volume of 200 µl within a plate well).

392

393 **Improvements in the application of the d-ROMs assay**

394 The d-ROMs assay can be performed using either cuvettes or well plates. The use of well  
395 plates may face the experimenter with an issue when there is any formation of precipitate  
396 on the bottom of the well. To overcome this problem, incubation can be done in tubes and,  
397 straightaway the end of incubation, tubes can be centrifuged and the supernatant used for  
398 the readings. Another problem may be with the plasma colour when this is very yellow. In  
399 this case, it is important to increase the wavelength at which readings are taken (e.g., 505  
400 nm or even more), considering that, for example, one peak of absorbance of lutein is at  $\approx$   
401 476 nm. As with the plasma colour, it should also be paid attention to whether plasma  
402 samples look red because of haemolysis. Iamele et al. (2002) found that a concentration of  
403 haemoglobin above 0.068 mM in human serum may interfere with the d-ROMs  
404 measurements.

405 The d-ROMs assay may not be sensitive, especially for young individuals, whose d-  
406 ROMs values may be low (Costantini et al. 2006). In order to improve the performance of  
407 the assay, it can be used more plasma than usual, but carefulness is needed in order to avoid  
408 any interference of plasma colour or alterations of the pH of the buffer.

409 Finally, it has been suggested that it might be interesting to measure fatty acids  
410 and/or cholesterol in plasma and to express d-ROMs values also per unit of fatty acids  
411 (Pérez-Rodríguez et al. 2015). This is because fatty acids and cholesterol are substrates for  
412 formation of organic hydroperoxides. Note that caution is needed because d-ROMs values  
413 do not only refer to lipid hydroperoxides but also to those formed from oxidation of other  
414 substrates and the correlations between d-ROMs values and either fatty acids or cholesterol  
415 are often low (e.g., Casagrande et al. 2011; Kotani et al. 2013). It might, however, be

416 interesting to assess if, together with uncorrected values, d-ROMs values corrected for the  
417 number of substrates that occur in plasma also provide valuable biological information.

418

## 419 **Conclusions**

420 Conversely to many colorimetric assays, the d-ROMs test has been properly assessed using  
421 electron paramagnetic resonance. All the direct and indirect evidence shows that the d-  
422 ROMs test is a valuable assay for the quantification of the plasma or serum oxidative status  
423 (higher values indicating higher oxidative damage), due to primary oxidation products of  
424 biomolecules (mainly organic hydroperoxides, but also endoperoxides and organic  
425 chloramines). Moreover, values of d-ROMs show significant individual repeatability over  
426 time (e.g., Costantini et al. 2007; Hau et al. 2015; Herborn et al. 2016). Experimental data  
427 also show that this assay is very sensitive to specific pro-oxidant agents and can provide  
428 ecologically valuable information, suggesting that this test might serve as a blood-derived  
429 biomarker to assess the impact of oxidative damage on health and fitness perspectives in  
430 animals. Clearly, as for any other metric of oxidative damage, the d-ROMs test should also  
431 be used in combination with other assays in order to better assess the individual oxidative  
432 balance. It is recommended that ecologists, interested in pursuing research in oxidative  
433 stress ecology, get sufficient basic background in biochemistry before making their own  
434 choice about what assay is suitable for their work.

435

## 436 **Acknowledgments**

437 I thank the Associate Editor, three anonymous reviewers, Angelo Alberti and Stefania  
438 Casagrande for providing valuable comments that helped me to improve the presentation

439 of the article; Katherine Herborn for kindly providing data on *Phalacrocorax aristotelis*  
440 reported in Table 3. Research Foundation of Flanders has supported this work.

441

## 442 **References**

443 Alberti A, Bolognini L, Macciantelli D, Caratelli M (2000) The radical cation of N,N-  
444 diethyl-para-phenyldiamine: a possible indicator of oxidative stress in biological  
445 samples. *Res Chem Intermediat* 26:253-267

446 Al-Johany AM, Haffor AS (2009) Effects of cadmium exposure on the ultrastructural  
447 pathology of different pulmonary cells, leukocyte count, and activity of glutathione  
448 peroxidase and lactate dehydrogenase in relation to free radical production in *Uromastyx*  
449 *aegyptius*. *Ultrastruct Pathol* 33:39-347

450 Assenza G, Zappasodi F, Squitti R et al (2009) Neuronal functionality assessed by  
451 magnetoencephalography is related to oxidative stress system in acute ischemic stroke.  
452 *NeuroImage* 44:1267-1273

453 Balkaya M, Prinz V, Custodis F et al (2011) Stress worsens endothelial function and  
454 ischemic stroke via glucocorticoids. *Stroke* 42:3258-3264

455 Banfi G, Malavazos A, Iorio EL, Dolci A, Doneda L, Verna R, Corsi MM (2006) The iron-  
456 *o*-dianisidine/xylenol orange assay in comparative oxidative stress assessment. Some  
457 possible shortcomings. *Eur J Appl Physiol* 97:506–508

458 Beaulieu M, González-Acuña D, Thierry AM, Polito MJ (2015) Relationships between  
459 isotopic values and oxidative status: insights from populations of gentoo penguins.  
460 *Oecologia* 177:1211-1220

461 Beaulieu M, Mboumba S, Willaume E, Kappeler PM, Charpentier MJ (2014) The oxidative  
462 cost of unstable social dominance. *J Exp Biol* 217:2629-2632

463 Bednarek J, Wysocki H, Sowinski J (2004) Oxidation products and antioxidant markers in  
464 plasma of patients with Graves' disease and toxic multinodular goiter: effect of  
465 methimazole treatment. *Free Radical Res* 38:659-664

466 Benedetti S, Lamorgese A, Piersantelli M, Pagliarani S, Benvenuti F, Canestrari F (2004)  
467 Oxidative stress and antioxidant status in patients undergoing prolonged exposure to  
468 hyperbaric oxygen. *Clin Biochem* 37:312-317

469 Bernabucci U, Colavecchia L, Danieli PP, Basiricò L, Lacetera N, Nardone A, Ronchi B  
470 (2011) Aflatoxin B1 and fumonisin B1 affect the oxidative status of bovine peripheral  
471 blood mononuclear cells. *Toxicol In Vitro* 25:684-691

472 Bertuglia S, Giusti A (2003) Microvascular oxygenation, oxidative stress, NO suppression  
473 and superoxide dismutase during postischemic reperfusion. *Am J Physiol Heart Circ*  
474 *Physiol* 285:H1064-1071

475 Blount JD, Vitikainen EIK, Stott I, Cant MA (2015) Oxidative shielding and the cost of  
476 reproduction. *Biol Rev* (published online, doi: 10.1111/brv.12179)

477 Boncoraglio G, Caprioli M, Saino N (2012) Solicitation displays reliably reflect oxidative  
478 damage in barn swallow nestlings. *Behav Ecol Sociobiol* 66:539–546

479 Bonisoli-Alquati A, Mousseau TA, Møller AP, Caprioli M, Saino N (2010) Increased  
480 oxidative stress in barn swallows from the Chernobyl region. *Comp Biochem Physiol A*  
481 155:205–210

482 Brambilla G, Civitareale C, Ballerini A, Fiori M, Amadori M, Archetti LI, Regini M, Betti  
483 M (2002) Response to oxidative stress as a welfare parameter in swine. *Redox Rep*  
484 7:159-163

485 Buonocore G, Perrone S, Longini M, Terzuoli L, Bracci R (2000) Total hydroperoxide and  
486 advanced oxidation protein products in preterm hypoxic babies. *Pediatr Res* 47:221-224

487 Cadet J, Di Mascio P (2006) Peroxides in biological system. In: Rappaport Z (ed) *The*  
488 *Chemistry of Peroxides*. John Wiley & Sons Ltd, Chichester, pp 915-999

489 Calabrese L, Carbonaro M (1986) An e.p.r. study of the non-equivalence of the copper of  
490 caeruloplasmin. *Biochem J* 238:291-295

491 Casagrande S, Costantini D, Groothuis TGG (2012) Interaction between sexual steroids and  
492 immune response in affecting oxidative status of birds. *Comp Biochem Physiol A*  
493 163:296–301

494 Casagrande S, Dell’Omo G, Costantini D, Tagliavini J, Groothuis T (2011) Variation of a  
495 carotenoid-based trait in relation to oxidative stress and endocrine status during the  
496 breeding season in the Eurasian kestrel: a multi-factorial study. *Comp Biochem Physiol*  
497 *A* 160:16-26

498 Casagrande S, Pinxten R, Zaid E, Eens M (2014) Carotenoids, birdsong and oxidative  
499 status: administration of dietary lutein is associated with an increase in song rate and  
500 circulating antioxidants (albumin and cholesterol) and a decrease in oxidative damage.  
501 *PLoS ONE* 9:e115899

502 Castellini C, Lattaioli P, Dal Bosco A, Minelli A, Mugnai C (2003) Oxidative status and  
503 semen characteristics of rabbit buck as affected by dietary vitamin E, C and n-3 fatty  
504 acids. *Reprod Nutr Dev* 43:91-103

505 Colombini F, Carratelli M, Alberti A (2016) Oxidative stress, d-ROMs test and  
506 ceruloplasmin. Free Radical Res (published online, doi:  
507 10.3109/10715762.2015.1136063)

508 Costantini D (2008) Oxidative stress in ecology and evolution: lessons from avian studies.  
509 Ecol Lett 11:1238-1251

510 Costantini D (2014) Oxidative stress and hormesis in evolutionary ecology and physiology:  
511 a marriage between mechanistic and evolutionary approaches. Springer-Verlag, Berlin  
512 Heidelberg

513 Costantini D, Casagrande S, De Filippis S, Brambilla G, Fanfani A, Tagliavini J, Dell’Omo  
514 G (2006) Correlates of oxidative stress in wild kestrel nestlings (*Falco tinnunculus*). J  
515 Comp Physiol B 176:329-337

516 Costantini D, Casasole G, AbdElgawad H, Asard H, Eens M (2016) Experimental evidence  
517 that oxidative stress influences reproductive decisions. Funct Ecol (published online,  
518 doi: 10.1111/1365-2435.12608)

519 Costantini D, Casasole G, Eens M (2014a) Does reproduction protect against oxidative  
520 stress? J Exp Biol 217:4237-4243

521 Costantini D, Coluzza C, Fanfani A, Dell’Omo G (2007) Effects of carotenoid  
522 supplementation on colour expression, oxidative stress and body mass in rehabilitated  
523 captive adult kestrels (*Falco tinnunculus*). J Comp Physiol B 177:723-731

524 Costantini D, Dell’Omo G (2006) Effects of T-cell-mediated immune response on avian  
525 oxidative stress. Comp Biochem Physiol A 145:137–142

526 Costantini D, Dell’Omo G (2015) Oxidative stress predicts long-term resight probability  
527 and reproductive success in the Scopoli’s shearwater (*Calonectris diomedea*). *Conserv*  
528 *Physiol* 3:cov024

529 Costantini D, Fanfani A, Dell’Omo G (2008) Effects of corticosteroids on oxidative damage  
530 and circulating carotenoids in captive adult kestrels (*Falco tinnunculus*). *J Comp Physiol*  
531 *B* 178:829-835

532 Costantini D, Ferrari C, Pasquaretta C, Cavallone E, Carere C, von Hardenberg A, Réale D  
533 (2012b) Interplay between plasma oxidative status, cortisol and coping styles in wild  
534 alpine marmots, *Marmota marmota*. *J Exp Biol* 215:374–383

535 Costantini D, Greives TJ, Hau M, Partecke J (2014b) Does urban life change blood  
536 oxidative status in birds? *J Exp Biol* 217:2994-2997

537 Costantini D, Monaghan P, Metcalfe NB (2011) Biochemical integration of blood redox  
538 state in captive zebra finches (*Taeniopygia guttata*). *J Exp Biol* 214:1148-1152

539 Costantini D, Monaghan P, Metcalfe N (2012a) Early life experience primes resistance to  
540 oxidative stress. *J Exp Biol* 215:2820–2826

541 Criscuolo F, Font-Sala C, Bouillaud F, Poulin N, Trabalon M (2010) Increased ROS  
542 production: a component of the longevity equation in the male Mygalomorph,  
543 *Brachypelma albopilosa*. *PLoS ONE* 5:e13104

544 Dotan Y, Lichtenberg D, Pinchuk I (2004) Lipid peroxidation cannot be used as a universal  
545 criterion of oxidative stress. *Progr Lipid Res* 43:200–227

546 Dudek K, Bednarek D, Siwicki AK, Rokita E, Studziński T (2013) The effect of LPS  
547 injections on non-specific immune response in affected pigeons. *Polish J Vet Sci*  
548 16:723-729

549 Ehrenwald E, Chisolm GM, Fox PL (1994) Intact human ceruloplasmin oxidatively  
550 modifies low density lipoprotein. *J Clin Invest* 93:1493-1501

551 Erel O (2005) A new automated colorimetric method for measuring total oxidant status.  
552 *Clin Biochem* 38:1103-1111

553 Favilli F, Marraccini P, Iantomasi T, Vincenzini MT (1997) Effect of orally administered  
554 glutathione on glutathione levels in some organs of rats: role of specific transporters.  
555 *Brit J Nutr* 78:293-300

556 Feng J, Ma WQ, Gu ZL, Wang YZ, Liu JX (2007) Effects of dietary copper (II) sulfate and  
557 copper proteinate on performance and blood indexes of copper status in growing pigs.  
558 *Biol Trace Elem Res* 120:171-178

559 Ferretti G, Bacchetti T, Principi F, Di Ludovico F, Viti B, Angeleri VA, Danni M,  
560 Provinciali L (2005) Increased levels of lipid hydroperoxides in plasma of patients with  
561 multiple sclerosis: a relationship with paraoxonase activity. *Mult Scler* 11:677-682

562 Fleming JJ, Santhosh S, Selvakumar R, Jose A, Eapen CE (2009) Usefulness of ferroxidase  
563 activity of ceruloplasmin in the diagnosis of Wilson's disease. *Indian J Clin Biochem*  
564 24:15-22

565 Geiger S, Le Vaillant M, Lebard T, Reichert S, Stier A, Le Maho Y, Criscuolo F (2012)  
566 Catching-up but telomere loss: half-opening the black box of growth and ageing trade-  
567 off in wild king penguin chicks. *Mol Ecol* 21:1500-1510

568 Gerardi G, Usberti M, Martini G, Albertini A, Sugherini L, Pompella A, Di LD (2002)  
569 Plasma total antioxidant capacity in hemodialyzed patients and its relationships to other  
570 biomarkers of oxidative stress and lipid peroxidation. *Clin Chem Lab Med* 40:104-110

571 Girotti AW (1998) Lipid hydroperoxide generation, turnover, and effector action in  
572 biological systems. *J Lipid Res* 39:1529-1542

573 Goldstein IM, Kaplan B (1979) Ceruloplasmin. A scavenger of superoxide anion radicals. *J*  
574 *Biol Chem* 254:4040–4045

575 Griffith OW, Meister A (1979) Potent and specific inhibition of glutathione synthesis by  
576 buthionine sulfoximine (S-n-butyl homocysteine sulfoximine). *J Biol Chem* 254:7558–  
577 7560

578 Halliwell BH, Gutteridge JMC (2007) *Free Radicals in Biology and Medicine*, 4th edn.  
579 Oxford University Press, Oxford

580 Harma MI, Harma M, Erel O (2006) Measuring plasma oxidative stress biomarkers in sport  
581 medicine. *Eur J Appl Physiol* 97:505

582 Hau M, Haussmann MF, Greives TJ, Matlack C, Costantini D, Quetting M, Adelman JS,  
583 Miranda AC, Partecke J (2015) Repeated stressors in adulthood increase the rate of  
584 biological ageing. *Front Zool* 12:4

585 Haussmann MF, Longenecker AS, Marchetto NM, Juliano SA, Bowden RM (2012)  
586 Embryonic exposure to corticosterone modifies the juvenile stress response, oxidative  
587 stress and telomere length. *Proc R Soc Lond B* 279:1447-1456

588 Herborn KA, Coffey J, Larcombe SD, Alexander L, Arnold KE (2011) Oxidative profile  
589 varies with personality in European greenfinches. *J Exp Biol* 214:1732–1739

590 Herborn KA, Daunt F., Heidinger BJ, Granroth-Wilding HMV, Burthe SJ, Newell MA,  
591 Monaghan P (2016) Age, oxidative stress exposure and fitness in a long-lived seabird.  
592 *Funct Ecol* (published online, doi: 10.1111/1365-2435.12578)

593 Hussain I, Zhang Y, Huang S (2014) Degradation of aniline with zero-valent iron as an  
594 activator of persulfate in aqueous solution. *RSC Adv* 4:3502

595 Isaksson C, Sepil I, Baramidze V, Sheldon BC (2013) Explaining variance of avian malaria  
596 infection in the wild: the importance of host density, habitat, individual life-history and  
597 oxidative stress. *BMC Ecol* 13:15

598 Ito Y, Nakamura A, Adachi M, Kato A, Ishikawa H, Maruta K, Hata T (2009) Blood  
599 oxidative stress levels and urinary dopamine values reflect the impact of academic stress  
600 on rodents. *J Anal Bio-Sci* 32:422-426

601 Jones DP (2006) Redefining oxidative stress. *Antioxid Redox Sign* 8:1865-1879

602 Kagan VE, Borisenko GG, Tyurina YY, Tyurin VA, Jiang J, Potapovich AI, Kini V,  
603 Amoscato AA, Fujii Y (2004) Oxidative lipidomics of apoptosis: redox catalytic  
604 interactions of cytochrome c with cardiolipin and phosphatidylserine. *Free Radical Biol*  
605 *Med* 37:1963

606 Kautz L, Jung G, Valore EV, Rivella S, Nemeth E, Ganz T (2014) Identification of  
607 erythroferrone as an erythroid regulator of iron metabolism. *Nat Genet* 46:678-684

608 Kilk K, Meitern R, Härmson O, Soomets U, Hõrak P (2014) Assessment of oxidative stress  
609 in serum by d-ROMs test. *Free Radical Res* 48:883-889

610 Kotani K, Tsuzaki K, Taniguchi N, Sakane N (2013) Correlation between reactive oxygen  
611 metabolites & atherosclerotic risk factors in patients with type 2 diabetes mellitus.  
612 *Indian J Med Res* 137:742-748

613 Kulmacz R, Miller JJ, Pendleton R, Lands W (1990) Cyclooxygenase initiation assay for  
614 hydroperoxides. *Methods Enzymol* 186:431-438

615 Kusuma Kumari M, Sankaranarayana T (2014) Evaluation of oxidative stress in type 2  
616 diabetes mellitus patients. IOSR J Dent Med Sci 13:46-50

617 Lajtha A, Tettamanti G, Goracci G (2009) Handbook of Neurochemistry and Molecular  
618 Neurobiology. Neural Lipids, 3rd edn. Springer Science, New York

619 Lee C, Yoon J (2004) Temperature dependence of hydroxyl radical formation in the  
620  $h\nu/Fe^{3+}/H_2O_2$  and  $Fe^{3+}/H_2O_2$  systems. Chemosphere 56:923-934

621 Liang Y, Roede JR, Dikalov S, Miller NG, Dudley Jr, SC, Quyyumi A, Jones DP (2012)  
622 Determination of ebselen-sensitive reactive oxygen metabolites (ebROM) in human  
623 serum based upon N,N'-diethyl-1,4-phenylenediamine oxidation. Clin Chim Acta 414:1-  
624 6

625 Lin H, Deculypere E, Buyse J (2004) Oxidative stress induced by corticosterone  
626 administration in broiler chickens (*Gallus gallus domesticus*) - 2. Short-term effect.  
627 Comp Biochem Physiol B 139:745-751

628 Lubrano V, Vassalle C, L'Abbate A, Zucchelli GC (2002) A new method to evaluate  
629 oxidative stress in humans. Immuno-Anal Biol Spe 17:172-175

630 Lucas LD, French SS (2012) Stress-induced tradeoffs in a free-living lizard across a  
631 variable landscape: consequences for individuals and populations. PLoS ONE 7:e49895

632 Lumeij JT, de Bruijne JJ (1985) Blood chemistry reference values in racing pigeons  
633 (*Columba livia domestica*). Avian Pathol 14:401-408

634 Marcus JR (1956) On the theory of oxidation-reduction reactions involving electron  
635 transfer. J Chem Phys 24:966

636 Maykova EV, Ismagilova RK, Alimova FK, Valeeva IH, Podolskaya AA, Kravtsova OA  
637 (2013) The association study of antioxidant status and antioxidant genes polymorphisms

638 in patients with ischemic heart disease in the Republic of Tatarstan. *World Appl Sci J*  
639 27:1367-1371

640 Metcalfe NB, Alonso-Alvarez C (2010) Oxidative stress as a life-history constraint: the role  
641 of reactive oxygen species in shaping phenotypes from conception to death. *Funct Ecol*  
642 24:984–996

643 Miyamoto S, Ronsein GE, Prado FM et al (2007) Biological hydroperoxides and singlet  
644 molecular oxygen generation. *IUCMB Life* 59:322-331

645 Miyazawa T (1989) Determination of phospholipid hydroperoxides in human blood plasma  
646 by a chemiluminescence-HPLC assay. *Free Radical Biol Med* 7:209-217

647 Montgomery MK, Buttemer WA, Hulbert AJ (2012) Does the oxidative stress theory of  
648 aging explain longevity differences in birds? II. Antioxidant systems and oxidative  
649 damage. *Exp Gerontol* 47:211-222

650 Montgomery MK, Hulbert AJ, Buttemer WA (2011) The long life of birds: the rat-pigeon  
651 comparison revisited. *PLoS ONE* 6:e24138

652 Musci G, Bellenchi GC, Calabrese L (1999) The multifunctional oxidase activity of  
653 ceruloplasmin as revealed by anion binding studies. *Eur J Biochem* 265:589-597

654 Nagatomo F, Fujino H, Kondo H, Ishihara A (2012) Oxygen concentration-dependent  
655 oxidative stress levels in rats. *Oxid Med Cell Longev* 2012:381763

656 Neyens E, Baeyens J (2003) A review of classic Fenton's peroxidation as an advanced  
657 oxidation technique. *J Hazard Mater* 98:33-50

658 Noguera JC, Lores M, Alonso-Alvarez C, Velando A (2011) Thrifty development: early-  
659 life diet restriction reduces oxidative damage during later growth. *Funct Ecol* 25:1144–  
660 1153

661 Ovrebø KK, Sørbye H, Svardal A, Grong K, Svanes K (1997) Glutathione and N-  
662 acetylcysteine reduce gastric mucosal blood flow in rats. *Digest Dis Sci* 42:1765-1774

663 Ovrebø KK, Svardal A (2000) The effect of glutathione modulation on the concentration of  
664 homocysteine in plasma of rats. *Pharmacol Toxicol* 87:103-107

665 Pasquini A, Luchetti E, Marchetti V, Cardini G, Iorio EL (2008) Analytical performances  
666 of d-ROMs test and BAP test in canine plasma. Definition of the normal range in  
667 healthy Labrador dogs. *Vet Res Commun* 32:137-143

668 Pérez-Rodríguez L, Romero-Haro AA, Sternalski A, Muriel J, Mougeot F, Gil D, Alonso-  
669 Alvarez C (2015) Measuring oxidative stress: the confounding effect of lipid  
670 concentration in measures of lipid peroxidation. *Physiol Biochem Zool* 88:345-351

671 Riedel W1, Lang U, Oetjen U, Schlapp U, Shibata M (2003) Inhibition of oxygen radical  
672 formation by methylene blue, aspirin, or alpha-lipoic acid, prevents bacterial-  
673 lipopolysaccharide-induced fever. *Mol Cell Biochem* 247:83-94

674 Samokyszyn VM, Miller DM, Reif DW, Aust SD (1989) Inhibition of superoxide and  
675 ferritin-dependent lipid peroxidation by ceruloplasmin. *J Biol Chem* 264:21-26

676 Saravanan G, Ponmurugan P (2013) S-allylcysteine improves streptozotocin-induced  
677 alterations of blood glucose, liver cytochrome P450 2E1, plasma antioxidant system, and  
678 adipocytes hormones in diabetic rats. *Int J Endocrinol Metab* 11:e10927

679 Sato H, Takahashi T, Sumitani K, Takatsu H, Urano S (2010) Glucocorticoid generates  
680 ROS to induce oxidative injury in the hippocampus, leading to impairment of cognitive  
681 function of rats. *J Clin Biochem Nutr* 47:224-232

682 Schneeberger K, Czirjak GA, Voigt CC (2013) Inflammatory challenge increases measures  
683 of oxidative stress in a free-ranging, long-lived mammal. *J Exp Biol* 216:4514-4519

684 Schneeberger K, Czirjak G, Voigt CC (2014) Frugivory is associated with low measures of  
685 plasma oxidative stress and high antioxidant concentration in free-ranging bats.  
686 *Naturwissenschaften* 101:285-290

687 Schöttker B, Brenner H, Jansen EH et al (2015a) Evidence for the free radical/oxidative  
688 stress theory of ageing from the CHANCES consortium: a meta-analysis of individual  
689 participant data. *BMC Med* 13:300

690 Schosinsky KH, Lehmann HP, Beeler MF (1974) Measurement of ceruloplasmin from its  
691 oxidase activity in serum by use of o-dianisidine dihydrochloride. *Clin Chem* 20:1556-  
692 63

693 Schöttker B, Saum KU, Jansen EH, Boffetta P, Trichopoulou A, Holleczeck B, Dieffenbach  
694 AK, Brenner H (2015b) Oxidative stress markers and all-cause mortality at older age: a  
695 population-based cohort study. *J Gerontol A Biol Sci Med Sci* 70:518-524

696 Sgorbini M, Bonelli F, Rota A, Marmorini P, Biagi G, Corazza M, Pasquini A (2015)  
697 Maternal and neonatal evaluation of derivated reactive oxygen metabolites (d-ROMs)  
698 and biological antioxidant potential in the horse. *Theriogen* 83:48-51

699 Sies H (1991) *Oxidative stress: oxidants and antioxidants*. London Academic Press,  
700 London

701 Simpson JA, Narita S, Giese S, Gebicki S, Gebicki JM, Dean RT (1992) Long-lived  
702 reactive species on free-radical damaged proteins. *Biochem J* 282:621-624

703 Sirak AA, Beavis AJ, Robertson FM (1991) Enhanced hydroperoxide production by  
704 peripheral blood leukocytes following exposure of murine epidermis to 12-  
705 Tetradecanoylphorbol-13-acetate. *Carcinogen* 12:91-95

706 Spolders M, Höltersshinken M, Meyer U, Rehage J, Flachowsky G (2010) Assessment of  
707 reference values for copper and zinc in blood serum of first and second lactating dairy  
708 cows. *Vet Med Int* 2010:194656

709 Stier A, Massemin S, Zahn S, Tissier ML, Criscuolo F (2015) Starting with a handicap:  
710 effects of asynchronous hatching on growth rate, oxidative stress and telomere dynamics  
711 in free-living great tits. *Oecologia* 179:999-1010

712 Suominen P, Punnonen K, Rajamäki A, Irjala K (1998) Serum transferrin receptor and  
713 transferrin receptor-ferritin index identify healthy subjects with subclinical iron deficits.  
714 *Blood* 92:2934-2939

715 Talukder S, Kerrisk KL, Gabai G, Fukutomi C (2015) Changes in milk oxidative stress  
716 biomarkers in lactating dairy cows with ovulatory and an-ovulatory oestrous cycles.  
717 *Animal Reprod Sci* 158:86-95

718 Talukder S, Kerrisk KL, Ingenhoff L, Gabai G, Garcia S, Celi P (2014) Changes in plasma  
719 oxidative stress biomarkers in dairy cows after oestrus synchronisation with controlled  
720 internal drug release (CIDR) and prostaglandinF<sub>2α</sub> (PGF<sub>2α</sub>). *Anim Prod Sci* 54:1490-  
721 1496

722 Tobler M, Ballen C, Healey M, Wilson M, Olsson M (2015) Oxidant trade-offs in  
723 immunity: an experimental test in a lizard. *PLoS ONE* 10:e0126155

724 Tomasetti M, Strafella E, Staffolani S, Santarelli L, Neuzil J, Guerrieri R (2010) α-  
725 tocopheryl succinate promotes selective cell death induced by vitamin K3 in  
726 combination with ascorbate. *Brit J Cancer* 102:1224-1234

727 van de Crommenacker J (2011) Hard times in paradise? Oxidative status, physiology and  
728 fitness in the tropical Seychelles warbler. PhD Thesis, University of Groningen

729 van de Crommenacker J, Komdeur J, Burke T, Richardson DS (2011a) Spatio-temporal  
730 variation in territory quality and oxidative status: a natural experiment in the Seychelles  
731 warbler (*Acrocephalus sechellensis*). *J Anim Ecol* 80:668–680

732 van de Crommenacker J, Komdeur J, Richardson DS (2011b) Assessing the cost of helping:  
733 the roles of body condition and oxidative balance in the Seychelles warbler  
734 (*Acrocephalus sechellensis*). *PLoS ONE* 6:e26423

735 Vassalle C, Boni C, Di Cecco P, Ndreu R, Zucchelli GC (2006) Automation and validation  
736 of a fast method for the assessment of in vivo oxidative stress levels. *Clin Chem Lab*  
737 *Med* 44:1372-1375

738 Wang J, Li D, Che L, Lin Y, Fang Z, Xu S, Wu D (2014) Influence of organic iron  
739 complex on sow reproductive performance and iron status of nursing pigs. *Livest Sci*  
740 160:89-96

741

742 Table 1 Examples of how changes of d-ROMs values in relation to a number of behavioural  
 743 traits or stressors have been found across a broad variety of organisms

<b>Trait or stressor</b>	<b>Taxon</b>	<b>Effect</b>	<b>Study</b>
Dominance status during the breeding season	<i>Acrocephalus sechellensis</i>	Dominant males had higher d-ROMs values	van de Crommenacker et al. 2011b
Dominance status during the breeding season	<i>Mandrillus sphinx</i>	High-ranking males had higher d-ROMs values	Beaulieu et al. 2014
Food habits	33 bat species	Species with a frugivorous diet had the lowest d-ROMs values, followed by omnivorous and animalivorous species	Schneeberger et al. 2014
Food habits	<i>Pygoscelis papua</i>	Colonies with the highest $\delta^{13}C$ and $\delta^{15}N$ values had the highest d-ROMs values	Beaulieu et al. 2015
Heat stress	<i>Taeniopygia guttata</i>	Increase of d-ROMs values	Costantini et al. 2012a
Immune response	<i>Falco tinnunculus</i>	Increase of d-ROMs values	Costantini and Dell'Omo 2006
Immune response	<i>Carollia perspicillata</i>	Increase of d-ROMs values	Schneeberger et al. 2013
Malaria infection	<i>Parus major</i>	Increase of d-ROMs values	Isaksson et al. 2013
Solicitation display	<i>Hirundo rustica</i>	Begging bout duration was negatively predicted by d-ROMs values but only after food deprivation.	Boncoraglio et al. 2012
Song behaviour	<i>Sturnus vulgaris</i>	Decrease of song rate with increase of d-ROMs values	Casagrande et al. 2014

744

745 Table 2 Comparison of absorbance values between samples that were either not treated or  
 746 treated with sodium azide (inhibitor of ceruloplasmin activity). Affinity of sodium azide for  
 747 ceruloplasmin (and so its capacity to inhibit ceruloplasmin activity) varies across species,  
 748 but a concentration of 1 mM of sodium azide in a buffer with a 7.4 pH has been shown to  
 749 reduce ceruloplasmin activity in both humans and chickens (Musci et al. 1999). The  
 750 capacity of sodium azide to inhibit ceruloplasmin was also suggested to increase with the  
 751 decrease of pH (Musci et al. 1999). This is important because the buffer used for the d-  
 752 ROMs assay has a pH of 4.8, hence much lower than that used in Musci et al. (1999). Data  
 753 on *Phalacrocorax aristotelis* are from Herborn et al. 2016. Values are expressed as mean  $\pm$   
 754 standard deviation. Coefficient of variation refers to variation in absorbance values between  
 755 samples that were either not treated or treated with sodium azide. Difference in absorbance  
 756 value mean between samples that were either not treated or treated with sodium azide is  
 757 also shown as percentage

Species	No sodium azide	1 mM sodium azide	Coefficient of variation	Difference expressed as %
<i>Serinus canaria</i>	0.132 $\pm$ 0.015	0.135 $\pm$ 0.017	3.03 $\pm$ 1.75	+3.68
<i>Sturnus vulgaris</i>	0.158 $\pm$ 0.021	0.163 $\pm$ 0.013	4.81 $\pm$ 1.34	+2.55
<i>Taeniopygia guttata</i>	0.182 $\pm$ 0.053	0.186 $\pm$ 0.052	2.20 $\pm$ 1.30	+1.88
<i>Calonectris diomedea</i>	0.110 $\pm$ 0.009	0.114 $\pm$ 0.009	2.66 $\pm$ 2.50	+3.95
<i>Phalacrocorax</i>	0.183 $\pm$ 0.025	0.175 $\pm$ 0.019	4.90 $\pm$ 2.50	-4.37

<i>aristotelis</i>				
--------------------	--	--	--	--

758

759

760 Table 3 Percentage of decrease of absorbance at 24 and 4°C as compared to 37°C, which is  
 761 the temperature required by the d-ROMs protocol. Note that the among sample variation is  
 762 to be expected given that in plasma samples there are several groups of organic  
 763 hydroperoxides that contribute to the reading and these differ in concentration among  
 764 individuals. Moreover, in plasma samples there are other primary oxidative damage  
 765 molecules that contribute to the reading (e.g., organic chloramines that derive from  
 766 oxidation of proteic amine groups; endoperoxides; Alberti et al. 2000; Liang et al. 2012)

767

768

	% of decrease of absorbance at 24°C as compared to 37°C	% of decrease of absorbance at 4°C as compared to 37°C
Cumene hydroperoxide 1	39	62
Cumene hydroperoxide 2	41	61
Standard 1	46	71
Standard 2	50	69
Plasma 1	42	60
Plasma 2	34	54
Plasma 3	36	57
Plasma 4	38	56
Coefficient of variation (%)	12.9	9.9

769

770 **Figure captions**

771 Fig. 1 Scheme illustrating the generation and fate of lipid hydroperoxides in cell  
772 membranes. PHGPx = phospholipid glutathione peroxidase; PLA2 = phospholipase A2;  
773 LOOH = fatty acid hydroperoxides; GPx = glutathione peroxidase; LOO• = fatty acid  
774 peroxy radical; LO• = fatty acid alkoxy radical; <sup>1</sup>O<sub>2</sub> = singlet oxygen; HOCl =  
775 hypochlorous acid; ONOO<sup>-</sup> = peroxynitrite. Russell mechanism refers to the reaction  
776 between peroxy radicals which generates singlet oxygen. Reprinted from Miyamoto et al.  
777 (2007) with permission from John Wiley and Sons


778  
779 Fig. 2 (A) Generation of hydroperoxides in protein backbone mediated by •OH attack. (B)  
780 Amino acid hydroperoxides containing hydrogen- $\alpha$ . Valine hydroperoxide (Val-OOH),  
781 lysine hydroperoxide (Lys-OOH) and leucine hydroperoxide (Leu-OOH). Reprinted from  
782 Miyamoto et al. (2007) with permission from John Wiley and Sons

783  
784 Fig. 3 Structures of thymidine and cytidine hydroperoxides formed by reaction of  
785 thymidine and cytidine with •OH, respectively. Reprinted from Miyamoto et al. (2007)  
786 with permission from John Wiley and Sons

787  
788 Fig. 4 Pre- and post-manipulation levels of red blood cell reduced glutathione, plasma d-  
789 ROMs values and plasma ceruloplasmin of canaries in relation to injection of sulfoximine.  
790 The experimental treatment was able to decrease red blood cell concentration of reduced  
791 glutathione and to increase plasma d-ROMs values, while it did not affect activity of  
792 ceruloplasmin. Although glutathione concentration was quantified within red blood cells, it  
793 is well established that sulfoximine is a potent inhibitor of glutathione synthesis in many  
794 body compartments, plasma included (see text). Means that do not share the same letter are  
795 significantly different from each other (Tukey,  $P < 0.05$ ). Data are shown as mean  $\pm$   
796 standard error. Data of d-ROMs and reduced glutathione are reprinted from Costantini et al.  
797 (2016) with slight modifications with permission from John Wiley and Sons

798  
799 Fig. 5 Dose-response of the d-ROMs test to pure solutions of cumene hydroperoxide. The  
800 concentrations of cumene hydroperoxides used in this trial are in the range of circulating  
801 organic hydroperoxides that occur in vertebrates. The concentrations of cumene  
802 hydroperoxides refer to those of the original samples, i.e., before they are 100 fold diluted  
803 when added to the d-ROMs buffer

804





805

806

807 Figure 1

808





809

810

811 Figure 2

812




813

814

815 Figure 3

816




817

818

819 Figure 4

820



821

822

823 Figure 5

824